



Bacillus-Care

复合芽孢杆菌

包装 Packaging: 1 公斤/1kg

可湿性粉剂 Soluble powder

有效成分: 多种芽孢杆菌菌群

2x10¹⁰ 孢子/克

Active Ingredients: Spores of various types of
Bacillus sp 2x10¹⁰ cfu/g



KEN MICROBES BIOTECH

Produced by:

Ken Microbes Biotech Sdn Bhd

No. 7 Jalan Permata 9A/KS09, Taman
Perindustrian Air Hitam Phase 2, , 41200
Klang, Selangor

功能/作用/ FUNCTIONS

主治根部病害：软腐病、枯萎病、根腐病、等。对叶面病害也有一定的防治作用。

在作物根部形成优势菌群，达到防病、固氮、解磷、解钾的功效。有效改良土壤结构。

Treats root diseases such as bacterial rot, Fusarium wilt, Phytium and Phytophthora, etc.

It also has a good treatment effect on leaf diseases. It becomes the dominant microbe in the roots for disease prevention, nitrogen fixation, phosphorus & potassium decomposition. Improve soil structure effectively.

使用说 / HOW TO USE

叶面喷洒: 1: 600-1000 倍 (看情况而定, 病情严重可加大剂量), 每 7-10 天用一次。

Leaf spray: 1 : 600-1000 times (Depending on the conditions, the dosage can be increased in severe cases), use every 7-10 days.

土壤灌根: 1-2 公斤/英亩, 每个月施用一次 | Irrigation/fertigation: 1-2kg/acre, apply once a month.

需要早期防治。用量更高, 效果更理想, 也不会影响作物。

Early prevention is needed. The higher the dose, the better the effect, and will not affect the crop.

注意事项 / PRECAUTIONS

可配合其它生物农药/化学杀虫药使用, 提高防治效。

Can be used together with other biological pesticides/chemical insecticides to enhance control effects.

不可混合杀菌剂或消毒剂使用, 避免益菌被杀死。

Cannot be used with fungicides or insecticides to avoid killing the bacillus.

建议在无病/发病初期使用。

Recommended for use in the early stages of the disease.



软腐病 Root Rot



枯萎病 Wilt Blight



猝倒病 Damping



根腐病 Root Disease



Tricho-Care

复合木霉菌

包装 Packaging: 1 公斤/1kg

水溶性粉剂 Soluble powder

有效成分: 哈兹木霉菌, 绿色木霉菌
5x10⁹ 孢子/克

Active ingredients: Beneficial fungal spores
(Trichorma viride, Trichoderma harzianum)
5x10⁹ cfu/g)



KEN MICROBES BIOTECH

Produced by:

Ken Microbes Biotech Sdn Bhd
No. 7 Jalan Permata 9A/KS09, Taman
Perindustrian Air Hitam Phase 2, , 41200
Klang, Selangor

功能/作用/ FUNCTIONS

根部性根腐病、枯萎病、腐霉病、灰霉病和白粉病等。对叶面病害也有一定的防治作用。

在作物根部形成优势菌群, 达到防病、固氮、解磷、解钾的功效。有效改良土壤结构。

Treats root diseases such as bacterial rot, Fusarium wilt, Phytiom and Phytophthora, Downy, etc. Also has a good treatment effect on leaf diseases. It becomes the dominant microbe in the roots for disease prevention, nitrogen fixation, phosphorus & potassium decomposition. Improves soil structure effectively.

使用说 / HOW TO USE

叶面喷洒: 1: 600-1000 倍 (看情况而定, 病情严重可加大剂量), 每 7-10 天用一次。

Leaf spray: 1 : 600-1000 times (Depending on the conditions, the dosage can be increased in severe cases), use every 7-10 days.

土壤灌根: 1-2 公斤/英亩, 每个月施用一次 | Irrigation/fertigation: 1-2kg/acre, apply once a month.

需要早期防治。用量更高, 效果更理想, 也不会影响作物。

Early prevention is needed. The higher the dose, the better the effect, and will not affect the crop.

注意事项 / PRECAUTIONS

不可混合杀菌剂或消毒剂使用, 避免益菌被杀死。

Do not use with fungicides or disinfectants to avoid killing trichoderma.

避免高温下使用以免影响真菌活性。不怕高湿, 湿度越高效果越好。

Avoid use at high temperatures so that fungal activity is not affected. The higher the humidity, the better the effect.

建议在无病/发病初期使用。

Recommended for use in the early stages of the disease.



软腐病 Root Rot



灰霉病 Botrytis



猝倒病 Damping



白粉病 White Mildew Disease



Premium Booster Plus 农抗抗菌肽



包装 Packaging: 1 公斤/1kg

水溶性粉剂 Soluble powder

有效成分: 抗菌肽蛋白

Active ingredients: Antimicrobial peptide proteir

Produced by:

Ken Microbes Biotech Sdn Bhd

No. 7 Jalan Permata 9A/KS09, Taman Perindustrian Air Hitam Phase 2, , 41200 Klang, Selangor

功能/作用/ FUNCTIONS

抗菌肽, 来于有益微生物发酵产物, 无残留, 无公害。有抑制坏菌作用。抗菌作用非常快速。

Has a fast and effective antibacterial effect. Antimicrobial peptide from beneficial microbial fermentation.

No residue, no pollution.

广谱抗菌剂: 白粉病、炭疽病、根腐病、锈病, 灰霉病等

Broad control of various diseases such as: Powdery mildew, Anthracnose, Root rot, Rust disease, Gray mold.

使用说 / HOW TO USE

叶面喷洒: 1: 400-500 倍 (看情况而定, 病情严重可加大剂量), 每 7-10 天用一次

Leaf spray: 1 : 600-1000 times (Depending on the conditions, the dosage can be increased in severe cases), use every 7-10 days.

浇根部: 20-30 克/20 升水 直到土壤完全湿润为止

Root spray (soil): 20-30g : 20 liters of water until the soil is completely moist.

需要早期防治。用量更高, 效果更理想, 也不会影响作物

Early prevention is required. The higher the dosage, the better the effect, and will not affect the crop.

注意事项 / PRECAUTIONS

可以复配其他杀菌剂或消毒剂使用, 效果增效。

Can be used in combination with chemical fungicides or other disinfectants to enhance the effect.

避免和有益微生物混合使用。Avoid using with beneficial microbes.

不能与碱性农药混用、混放。

Cannot be mixed or used together with alkaline pesticides.



Rust disease 锈病



灰霉病 Botrytis



炭疽病 Anthracnose



露菌病 Downy Disease



PLANT PROTECT

树医生



KEN MICROBES BIOTECH

包装 Packaging: 1 公斤/1kg

水溶性粉剂 Soluble powder

有效成分: 稳定亚磷酸, 抗病因子

Active Ingredients: Stabilized Phosphorus acids,
Immune Boosters

Produced by:

Ken Microbes Biotech Sdn Bhd

No. 7 Jalan Permata 9A/KS09, Taman
Perindustrian Air Hitam Phase 2, , 41200
Klang, Selangor

功能/作用/ FUNCTIONS

树医生的抗病作用是一种后天获得的系统性抗病作用。作物施用后, 可被植物叶片、根系吸收, 运送至植株体内, 发挥其直接杀菌功能, 同时启动抗病防御系统, 待病原菌入侵植株时, 刺激植株产生植物防御素从而产生抗病能力。对溃疡病、疫病菌、霜霉菌和白粉病等真菌病害具有很好的抑制效果。

When the plant is treated with this Plant Protect, the 'Phosphorus acid' in it can be absorbed by the leaves and roots of the plant, transported throughout the plant, and shows its anti-fungal function directly. At the same time, it activates the anti-disease defense system. When a pathogen attacks a plant, it stimulates the plant to produce plant defense substances, giving the plant the ability to fight disease. It has a good inhibitory effect against diseases such as Canker (Phytophthora), Downy Mildew and Downy Powder, and so on.

使用说 / HOW TO USE

叶面喷洒: 1: 600-1000 倍 (看情况而定, 病情严重可加大剂量), 每 7-10 天用一次。

Foliar spray: 1: 600-1000 times (Depending on the situation, the dosage can be increased in severe cases), use every 7-10 days.

冲施每亩每次 3 公斤; For spraying, use 3kg for each acre once.

治疗根部病害, 稀释 200-300 倍灌根; For treating root diseases, dilute 200-300 times and spray to the roots.

注意事项 / PRECAUTIONS

树医生混配性好, 可以与铜制剂复配一起使用, 不会产生沉淀与药害, 但不能与强碱制剂使用;

Plant Protect has the ability to be mixed with various kinds of drugs, it can be mixed with copper products without producing any precipitate or damage from the drug; But it cannot be used together with poisons or strong alkaline fertilizers.

亚磷酸钾可与常见的中微量元素肥料 (糖醇钙、硝酸钙等) 任意混用, 使用方便;

Can be mixed with common microelement fertilizers (such as calcium sorbitol, calcium nitrate) easily, convenient to use.



白粉病
White Mildew Disease



露菌病
Downy Mildew Disease



溃疡病
Canker Disease



白粉病
White Mildew Disease



Bio F1

抗病专家

包装 Packaging: 1 公斤/1kg

水溶性粉剂 Soluble powder

活性成分: 木霉属、芽孢杆菌属

Active Ingredients: Trichoderma sp, Bacillus sp



KEN MICROBES BIOTECH

Produced by:

Ken Microbes Biotech Sdn Bhd

No. 7 Jalan Permata 9A/KS09, Taman

Perindustrian Air Hitam Phase 2, , 41200

Klang, Selangor

功能/作用/ FUNCTIONS

含有多钟有益微生物的生物防治剂。保护树根系统，抵抗土壤病原体带来的疾病。

Biocontrol agent containing a variety of beneficial microbes. Protects the root system of trees against diseases from soil pathogens.

广泛控制多种病害，例如：镰刀菌、菌核、丝核菌、疫霉。

Broad control for various diseases such as: Fusarium, Sclerotium, Rhizoctonia, Phytophthora.

使用说 / HOW TO USE

叶面肥: 1 公斤: 每 5-7 天 1000 升水。

Foliar: : 1kg : 1000 liters of water every 5-7 days.

塑料袋: 1 公斤: 5000 个塑料袋灌溉施肥时需要 1000 升水。

Polybag: 1kg : 1000 liters of water by fertigation for 5000 polybags.

需要早期防治。用量更高，效果更理想，也不会影响作物。

Early prevention is needed. Higher dosage, better effect, and will not affect the crop.

注意事项 / PRECAUTIONS

适合早上或晚上使用。

Suitable for use in the morning or evening.

避免与化学杀菌剂/杀虫剂同时使用。

Avoid use with chemical fungicides/pesticides.

必须用水稀释后食用。

Must be used after dilution with water.



黄叶病
Banana Panama Disease



露菌病
Downy Mildew Disease



青枯病
Bacterial Wilt



Farm Clear

抗菌剂

包装 Packaging: 1 升/1L

有效成分: 消毒剂、抗菌剂

Active Ingredients: Disinfection and anti-microbe agents



KEN MICROBES BIOTECH

Produced by:

Ken Microbes Biotech Sdn Bhd

No. 7 Jalan Permata 9A/KS09, Taman Perindustrian Air Hitam Phase 2, , 41200 Klang, Selangor

功能/作用/ FUNCTIONS

用于农业和园艺的广谱消毒和抗菌控制剂。

A broad spectrum disinfectant and antimicrobial control agent for agriculture and horticulture.

有效对抗病原真菌、细菌和病毒。也可用于杀灭藻类。

Effective against pathogenic fungi, bacteria and viruses. Can also be used to kill algae.

使用说 / HOW TO USE

叶面喷施: 1L: 1000 升水喷洒在叶面和叶下。

Foliar: 1L : 1000 Liters of water spray on the surface and carry leaves.

注意事项 / PRECAUTIONS

适合早上或晚上使用。

Suitable for use in the morning or evening.

避免与化学杀菌剂/杀虫剂同时使用。

Avoid use with chemical fungicides/pesticides.

必须用水稀释后食用。

Must be used after dilution with water.



炭疽病 Anthracnose



露菌病
Downy Mildew Disease



青苔 Algae